

## LAND O LAKES® NURSING FORMULA BOV FF – 50#

**SPECIE:** Calf **TYPE OF FEED:** Calf Milk Replacer – Growth **FORM OF FEED:** Powder

### GENERAL DESCRIPTION

LAND O LAKES® Nursing Formula is a growth calf milk replacer. See features/benefits list below for more information.

### MIXING & FEEDING INSTRUCTIONS

#### USE ONLY LOW SODIUM (<50 ppm) water for mixing and feeding!

- Always weigh milk replacer powder for accurate mixing.
- Mix milk replacer powder according to the schedule below by adding powder to water as indicated and mix thoroughly

|                 | Milk Replacer Powder | Water (110-120° F) |
|-----------------|----------------------|--------------------|
| Individual Calf | 0.75 lbs             | + 2 quarts         |
| Large Batch     | 1.5 lbs              | + 1 gallon         |

#### Feeding Directions (2 days to Weaning):

##### NOTE: Always offer free choice water

**Large Breeds (120 lb of body weight):** Feed 2.3 quarts twice daily (fill 2 quart bottle to rim).

Mixed and fed as above, this product provides 45 milligrams of lasalocid per 100 lb of body weight. This resulting mix should be fed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

**IMPORTANT:** Thoroughly mix milk replacer powder in warm water, and feed immediately. When preparing large batches (i.e., anything other than individual calf servings) keep the mixture agitated while dispensing into individual feeding containers to prevent any settling which could result in the feeding of improper levels of lasalocid.

#### General Recommendations:

1. Feed newborn calves 4 quarts of high quality warm colostrum via nipple bottle or esophageal feeder within 1 hour of birth and repeat 12 hours later via nipple bottle as much as they will consume.
2. Consult with your veterinarian on a dry cow and calf vaccination program including but not limited to E. Coli, rota and corona virus, clostridium and salmonella.
3. For best mixing, continuously stir with wire whip or power mixer while adding powder to 110-120° F. water. Use a thermometer to assure correct water temperature to avoid product separation and scalding. Feed milk replacer at 100- 105° F.
4. Begin feeding milk replacer on day 2 and provide fresh, clean low sodium (<50ppm) water along with a high quality, palatable calf starter on a free-choice basis.
5. At the start of weaning, transition calves from twice per day feeding to once per day feeding. This will encourage the calf to eat more calf starter and assures an effective weaning.
6. Wean only if calf is eating at least 3 lbs calf starter for 3 consecutive days.

**Ingredients:** Dried Whey Protein Concentrate, Dried Whey, Dried Whey Product, Dried Skimmed Milk, Dried Milk Protein, Animal Fat (preserved with BHA and BHT), Lecithin, Polysorbate 80, Dicalcium Phosphate, L-Lysine, DL-Methionine, L-Threonine, Calcium Carbonate, Yeast Extract, Vitamin A Acetate, Vitamin D3 Supplement, Vitamin E Supplement, Thiamine Mononitrate, Ascorbic Acid, Calcium Pantothenate, Niacin Supplement, Riboflavin, Biotin, Menadione Sodium Bisulfite Complex (source of Vitamin K3), Pyridoxine Hydrochloride, Folic Acid, Vitamin B12 Supplement, Choline Chloride, Calcium Silicate, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Ethylenediamine Dihydroiodide, Selenium Yeast and Natural & Artificial Flavor.

| GUARANTEED ANALYSIS           |              |
|-------------------------------|--------------|
| Crude Protein, not less than  | 22.0%        |
| Crude Fat, not less than      | 20.0%        |
| Crude Fiber, not more than    | 0.15%        |
| Calcium (Ca), not less than   | 0.75%        |
| Calcium (Ca), not more than   | 1.25%        |
| Phosphorus (P), not less than | 0.70%        |
| Vitamin A, not less than      | 20,000 IU/lb |
| Vitamin D3, not less than     | 5,000 IU/lb  |
| Vitamin E, not less than      | 150 IU/lb    |

**ACTIVE DRUG INGREDIENT** Lasalocid (as lasalocid sodium) . . . . . 72 gm/ton Equal to 36 mg/lb  
For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves.

**CAUTION:** The safety of lasalocid in unapproved species has not been established. This product is to be fed to calves only. Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

**WARNING:** A withdrawal period has not been established for lasalocid in pre-ruminating calves. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL. DO NOT FEED TO LACTATING COWS.

**FEATURES/  
BENEFITS:**

| FEATURES                  |  | BENEFITS                                                                                                                           |  |
|---------------------------|--|------------------------------------------------------------------------------------------------------------------------------------|--|
| <b>Prebiotic</b>          |  | Supports an optimal microbiome and immune health by feeding the good microorganisms                                                |  |
| <b>Amino Acid Balance</b> |  | Added amino acids such as methionine, threonine and lysine which are essential for optimal growth                                  |  |
| <b>Essential Oils</b>     |  | Helps to optimize digestibility and nutrient absorption; supports palatability which assists with milk replacer and starter intake |  |
| <b>Coccidiostat</b>       |  | Helps to control coccidiosis                                                                                                       |  |